

# Imported Molassed Sugar Beet Feed



A palatable high digestible fibre and sugar energy, pelleted feed.

#### Typical Analysis (on a dry matter basis)

Dry matter (%)	Energy (MJ ME/kg DM)	Crude protein (%)	Oil (%)	NDF (%)	Starch (%)	Sugar (%)	DUP %
91.0	12.5	9.0	1.1	32.0	1.0	15.6	2.9

### What are you trying to achieve?

Need	Feature	Benefit	
Drive intake	A highly palatable feed.	Can stimulate intakes of less palatable feeds, increasing milk and meat production.	
Increase milk fat %	A good source of digestible fibre.	Provides the building blocks for milk fat synthesis, increasing value per litre.	
Increase energy intakes	Good levels of non-starch digestible fibre energy.	Allows energy intakes to be increased without increasing the risk of acidosis associated with cereal feeding.	
Minimise risk of acidosis	High content of digestible fibre.	Assists in maintaining an optimum rumen pH.	
Feeding flexibility	12mm durable pellet. Suitable for a wide range of livestock.	Specifically designed for floor feeding but can also be used in automated feeding systems. Can be transferred to feeders via auger systems.	

The predicted responses (benefits) assume that the specified nutrient, physical or structural dietary components are limiting livestock performance in the current ration.



#### **Complementary Concentrate Feeds**

- High starch feeds e.g. cereals, maize meals, and confectionary and bakery products.
- **High protein feeds** e.g. soya bean meal, rapeseed meal, wheat distillers

#### Recommended daily feed rates (per head basis)

Imported Molassed Sugar Beet Feed can be fed via automated feeders, top dressed or floor fed, and used individually or as part of a blend or TMR

Milking Cows	Up to 6 (typically 3)kg
Dry Cows	Up to 2 kg
Replacement Heifers	Up to 2 kg and up to 40% of the DMI
Calves (to 12 weeks)	Up to 1.5 kg and up to 40% of the DMI
Growing Cattle	Up to 2.5 kg and up to 40% of the DMI
Finishing Cattle	Up to 5kg and up to 50% of the DMI
Suckler Cows	Up to 4 (typically 2)kg
Ewes and Rams	Up to 1 (typically 0.5) kg
Hoggets and Lambs	Up to 1 kg or up to 50% of the DMI

DMI = dry matter intake

#### Availability, handling and storage

Imported Molassed Sugar Beet Feed is available all year round, UK wide as bulk tipped or blown loads only. Like all dry feeds, they should be stored in a secure shed, bunker, bin or hopper and kept cool, dry and free from vermin. Imported Molassed Sugar Beet Feed must be used within 12 months of delivery.

#### **Additional information**

#### Method of production

Imported Molassed Sugar Beet Feed is a co-product from sugar production. Once the sugar has been diffused out from the beet, the fibrous residues are dried and combined with beet molasses and pelleted.

#### **Quality Assurance**

Imported Molassed Sugar Beet Feed is FEMAS assured (or a recognised equivalent). Imported Molassed Sugar Beet Feed (Dried (sugar) beet feed, molassed) are listed under number 4.1.11 in the EU Catalogue of Feed Materials.



# Imported Molassed Sugar Beet Feed

## **Detailed Typical Analysis** (fresh basis other than where stated)

Oil A % 0.50 Magnesium g/kg 1.30   Oil B % 1.10 Phosphorus g/kg 0.80   Crude protein % 8.20 Potassium g/kg 28.0   Crude protein: % 9.00 Salt g/kg 1.00   DM - 16.3 Sodium g/kg 0.20   Ash % 8.70 Copper mg/kg 7.00   ME* - in vivo MJ/kg DM 12.5 Manganese mg/kg 40.0   NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of CP	Dry matter	%	91.0	Calcium	g/kg	8.00
Oil B % 1.10 Phosphorus g/kg 0.80   Crude protein % 8.20 Potassium g/kg 28.0   Crude protein: % 9.00 Salt g/kg 1.00   DM B.70 Salt g/kg 0.20   Ash % 8.70 Copper mg/kg 7.00   ME* - in vivo MJ/kg DM 12.5 Manganese mg/kg 40.0   NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of cP 5.31   aDM 0.50 Methionine % of CP 1.52	•					
Crude protein % 8.20 Potassium g/kg 28.0   Crude protein: % 9.00 Salt g/kg 1.00   DM 16.3 Sodium g/kg 0.20   Ash % 8.70 Copper mg/kg 7.00   ME* - in vivo MJ/kg DM 12.5 Manganese mg/kg 40.0   NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.52   cDM				<u> </u>		
Crude protein: % 9.00 Salt g/kg 1.00   DM 16.3 Sodium g/kg 0.20   Ash % 8.70 Copper mg/kg 7.00   ME* – in vivo MJ/kg DM 12.5 Manganese mg/kg 40.0   NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12						
DM Fibre % 16.3 Sodium g/kg 0.20   Ash % 8.70 Copper mg/kg 7.00   ME* – in vivo MJ/kg DM 12.5 Manganese mg/kg 40.0   NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN <t< td=""><td></td><td></td><td></td><td></td><td></td><td></td></t<>						
Fibre % 16.3 Sodium g/kg 0.20   Ash % 8.70 Copper mg/kg 7.00   ME* – in vivo MJ/kg DM 12.5 Manganese mg/kg 40.0   NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.52   cDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 4.55   aN 0.32	•	%	9.00	Salt	g/kg	1.00
Ash % 8.70 Copper mg/kg 7.00   ME* - in vivo MJ/kg DM 12.5 Manganese mg/kg 40.0   NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.52   cDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 4.55   aN 0.32 Threonine % of CP 4.55   bN 0.63 0.63		<del></del>	1 / 0	0 "		
ME* - in vivo MJ/kg DM 12.5 Manganese mg/kg 40.0   NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32 Description % of CP 4.55						
NDF % 29.0 Selenium mg/kg 0.09   Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32 DN 0.63	Ash		8.70	Copper	mg/kg	7.00
Starch % 0.60 Zinc mg/kg 20.0   Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	ME* – in vivo	MJ/kg DM	12.5	Manganese	mg/kg	40.0
Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	NDF	%	29.0	Selenium	mg/kg	0.09
Sugar % 14.2 Saturates % of oil 23.0   ERDP-FiM* % @ 6% 4.00 Monounsaturates % of oil 11.0   DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	Starch	%	0.60	Zinc	mg/kg	20.0
DUP-FiM* % @ 6% 2.90 PUFAs % of oil 66.0   DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	Sugar	%	14.2	Saturates		23.0
DUP digestibility % 70.0 Long chain PUFAs % of oil 0.00   sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	ERDP-FiM*	% @ 6%	4.00	Monounsaturates	% of oil	11.0
sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	DUP-FiM*	% @ 6%	2.90	PUFAs	% of oil	66.0
sDM 0.20 Lysine % of CP 5.31   aDM 0.50 Methionine % of CP 1.26   bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	DUP digestibility	%	70.0	Long chain PUFAs	% of oil	0.00
bDM 0.40 Cysteine % of CP 1.52   cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63			0.20	Lysine	% of CP	5.31
cDM 0.11 Histidine % of CP 3.92   sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	aDM		0.50	Methionine	% of CP	1.26
sN 0.12 Threonine % of CP 4.55   aN 0.32   bN 0.63	bDM		0.40	Cysteine	% of CP	1.52
aN 0.32   bN 0.63	cDM		0.11	Histidine	% of CP	3.92
bN 0.63	sN		0.12	Threonine	% of CP	4.55
	aN	<u>-</u>	0.32	·	·	
cN 0.06	bN	·	0.63		·	
	cN	·	0.06			